

In the Claims:

Please amend the claims as set out below:

1(currently amended).                    A reagent for use in an assay of an analyte, said reagent comprising:

    a fluid medium containing;

    a first substrate having a first binding species attached thereto, said first binding species capable of dissociating from the first substrate; and

    a second substrate having binding regions said binding region having binding partners capable of selectively binding dissociated first binding species without detrimentally affecting the signal strength of said assay.

2 (previously presented).            The reagent of claim 1, wherein the first and the second substrates are formed of the same materials.

3 (previously presented).            The reagent of claim 1, wherein the first and the second substrates are formed of different materials.

4(previously presented).            The reagent of claim 1, wherein said first binding species is selected from the group consisting of biotin, avidin, streptavidin, an antigen, an antibody, a hapten, a receptor and an oligonucleotide.

5(previously presented)            The reagent of claim 4, wherein said binding partner selectively binds species selected from the group consisting of biotin, avidin, streptavidin, an antigen, an antibody, a hapten a receptor and an oligonucleotide.

6(original).                    The reagent of claim 1, wherein said binding regions are selected from the group consisting of pores, interstitial spaces, indentations and grooves.

7(currently amended). A reagent comprising:

(1) a fluid medium containing;

(2) including a binding species attached to a substrate said binding species capable of disassociating from the substrate;

(2) a non-porous material including (a) an outer surface porous to dissociated binding species but not to the binding species attached to the substrate and (b) an inner surface having an affinity for said binding species

8(previously presented). The reagent of claim 7, wherein the binding species is selected from the group consisting of biotin, avidin, streptavidin, an antigen, an antibody, a hapten, a receptor and an oligonucleotide.

9 (withdrawn) A method of stabilizing a reagent through the inactivation of a free species, comprising:

providing a reagent storage container including a liquid medium containing a binding species attached to a substrate;

a permeable material including an inner surface having an affinity for said binding species; and

an outer surface permeable to the binding species when dissociated from said substrate.

10(withdrawn) The reagent of claim 9, wherein the first binding species is attached to a defined binding surface on the first substrate, and the binding regions on the second substrate are formed on a defined binding surface on the second substrate, and wherein the defined binding surface on the first substrate and the defined binding surface on the second substrate are formed of the same materials

11(withdrawn). The method of claim 10, wherein the binding species is selected from the group essentially consisting of biotin, avidin, streptavidin, an antigen, an antibody, a hapten, a receptor and an oligonucleotide, and bindable derivatives thereof.

12(currently amended). A reagent, comprising:

- a fluid medium containing;
- a first substrate ~~in contact with the fluid medium~~;
- a first binding species, wherein a first portion of the first binding species is attached to the first substrate and a second portion of the first binding species is dissociated from the first substrate; and
- a second substrate having regions adapted to selectively bind the second portion of the first binding species.

13 (cancelled).

14(currently amended). A reagent comprising:

- a fluid medium containing;
- a first substrate having a first binding species attached thereto, said first binding species capable of dissociating from the first substrate; and
- a second substrate having regions capable of selectively binding said dissociated first binding species without detrimentally affecting the signal strength of said assay and second regions capable of selectively binding first binding species associated with said first substrate.

15(previously presented). The reagent of claim 14, wherein the first substrate and the second substrate are formed of the same materials

16(previously presented). The reagent of claim 14, the first substrate and the second substrate are formed of different materials.

17(withdrawn)      A method of reducing the signal to noise ratio amplifying a signal in an assay comprising:

    providing a fluid medium containing a first substrate having at least one first binding species; and

    providing a second substrate having regions characterized by their ability to bind said first binding species if dissociated from said first substrate and also to bind said first substrate having said first binding species attached.

18(withdrawn).      The method of claim 17, wherein said first substrate comprises a plurality of particles.

19(withdrawn).      The method of claim 17, wherein said regions bind said first binding species from a plurality of particles.

20 (withdrawn).      The method of claim 17, wherein said second substrate is selected from the group consisting of latex particles, proteins such as bovine serum albumin (BSA), human serum albumin (HSA), antibodies, or polymers such as poly-BSA, poly-antibodies, dextrans, and dendrimers

21(withdrawn)      The method of claim 20, wherein said regions characterized by their ability to selectively bind said first binding species if dissociated from said first substrate are adapted to exclude said first species associated with said first substrate